



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

NOTE ON PROF. NICHOLSON'S SINGULAR VALUE OF II.

BY PROF. WILLIAM WOOLSEY JOHNSON.

IF we regard equation (6), p. 150, as simply equivalent to  $\infty.0 = 2 \div \pi$  it presents no difficulty; but if, on the other hand, the symbols  $(1-1)^{\frac{1}{2}}$  and  $(1-1)^{-\frac{1}{2}}$  be supposed to stand for the limits of  $(1-x)^{\frac{1}{2}}$  and  $(1-x)^{-\frac{1}{2}}$  when  $x = 1$ , the result appears paradoxical, since then the product of these quantities and therefore its limit is equal to unity.

The former is in fact correct for although equation (2) is the result of putting  $x = 1$  in the expansion of  $(1-x)^n$ , (3) is not, since the process of transforming the infinite series into an infinite product is applicable only when  $x = 1$ . Thus equation (3) means nothing more than that the infinite product in the second member has zero for its limit; in like manner equation (4) means only that the infinite product in its second member has no limit.

Moreover, the product obtained by taking an infinite number of factors from each series may have any value we choose, for this value is a function of the ratio of the infinite numbers of the factors taken from the two series. If  $p$  factors be taken from (4) and  $q$  factors from (3) the value of the product, when  $p$  and  $q$  are both infinite but  $p \div q = a$ , is

$$\frac{a^n \sin n\pi}{n\pi};$$

putting  $n = \frac{1}{2}$  and assuming  $a = 1$ , as implied in the manner of writing Wallis' Theorem, the result becomes  $2 \div \pi$ .

NOTE ON EXPERIMENTAL CONFIRMATION OF THEORETICAL DEDUCTION, BY THE EDITOR.—IF a plane surface is ruled with parallel and equidistant lines and a slender rod, the length of which equals the perpendicular distance between two consecutive lines, is thrown at hazard upon the plane, the probability that it will fall across a line is  $2 \div \pi$ . (See *Mathematical Monthly*, Vol. II, p. 236.)

If we denote this probability by  $P$ , we shall have

$$P = \frac{2}{\pi} = \frac{2}{3.14159} = .6366.$$

Hence a rod thrown at hazard upon the plane 10,000 times should fall across a line 6366 times.

At the recent Montreal meeting of the American Association for the Advancement of Science, Prof. Mendenhall exhibited before Section A. of the Association the result of 30,000 experiments which he had performed by